

Overview of C-Farm® simulation model

Model Developers: Armen R. Kemanian (akemanian@brc.tamus.edu)

Claudio O. Stöckle (stockle@wsu.edu)

Programming Support: Shawn Quisenberry

C-Farm is a cropping systems model that simulates, in a daily time step, the soil carbon balance, water balance, crop growth and yield, and other processes. The main objective is to provide accurate simulations of soil carbon and nitrous oxide emissions when accumulated over 3 to 5 years. The inputs to the model are simple, readily accessible, or easy to define. The inputs to the model are: (1) a daily weather file that includes the location description (latitude, longitude, and elevation), (2) a soil description for the site, layer by layer, (3) a cropping sequence, (4) basic agronomic info about each crop (average planting, flowering, and maturity date), (5) a list of the tillage tools used (from a drop down menu of available tools) with the approximate date for each operation, (6) irrigation set up (automatic or fixed-date irrigation). Once the weather and soil files are available, it takes a few minutes to set up a simulation. The outputs are available in a daily or annual basis for several soil and crop variables, including crop yield, residues yield, root biomass per layer, soil carbon balance by layer, crop transpiration, soil evaporation, runoff, percolation, and other variable of interest. We are in the process of linking the carbon cycling with already developed nitrogen cycling algorithms to provide estimates of nitrous oxide emissions. The model input/output interface is user-friendly and accessible through Excel spreadsheets. A stand-alone version will substitute the current beta version in VBA.

The basic input information is relatively easy to acquire but it has to be accurate for the model to provide realistic estimates of carbon storage rates for a given site. In particular, the current soil carbon content needs to be provided as initial soil carbon is a key element affecting carbon storage rates.

Unique features of C-Farm

Three general features are unique in C-Farm:

(1) **Soil carbon cycling:** the carbon cycling subroutine has been originally developed for this model. Soil carbon is composed of a single carbon pool whose turnover rate depends on the size of the pool. The higher the carbon pool, the higher the turnover rate. Unlike C-Farm, in multi-compartment models the soil carbon pool is subdivided in several pools (typically three) each having a different turnover rate. Thus, the overall turnover rate depends on the relative amount of each pool in the total soil carbon. A weakness of the multi-pool model is the difficulty in setting up the actual turnover rate of each pool, and the implicit assumption that as a carbon pool increases or decreases in size, the turnover rate remains stable, which is likely not true. The approach taken in C-Farm to regulate the soil carbon turnover rate is much simpler, and empirical evidence suggests it is sufficient to represent carbon turnover for different soils and climates. Similarly, the humification rate (the fraction of the decomposed carbon in plant residues or manure that remains in the soil as humus after an “effective” microbial attack) is controlled by the soil carbon

level in each layer. We consider that physical protection exerts a major control on humification rate (Withmore and Hassink, 1997; Six et al, 20XX). Therefore, clay content and soil carbon content control the humification rate of each layer. These are, of course, empirical approaches to represent complex processes.

(2) **Tillage effect on carbon turnover:** Tillage accelerates the turnover rate of soil carbon; however, no robust and scientific accepted procedure has been developed to represent this effect. In C-Farm, we developed a methodology to accelerate the turnover rate which is based on soil texture, the tool performing the operation (based on information developed by USDA National Resource Conservation Service for different tools), and the number of operations.

(3) **Soil water balance:** there are multiple robust models for soil water balance. Nonetheless, and to conform to the structure of C-Farm and its computational efficiency, new sub-routines were developed for soil water redistribution (for water above field capacity) and soil water evaporation. These approaches accurately reproduce water flow in the profile and the typical two-stage soil water evaporation rate for soils with different texture, without resorting to a numerical solution to the water flow in soils.

(4) **Crop growth and radiation interception:** crop growth is driven by radiation or transpiration. The transpiration-driven growth in C-Farm follows the approach outlined in Kemanian et al. (2005), which introduced a variant to the Tanner (1981) and Tanner and Sinclair (1983) approach. This modification improved the simulation of biomass production and therefore the estimation of carbon inputs. To the best of our knowledge this is the first time it is parameterized for several crops and used in a crop simulation model.

Below are screen shots of the C-Farm model Excel Interface, which is a beta version written in Visual Basic for Applications. A standalone version in Visual Basic substituted this beta version.

Input Control Interface

Microsoft Excel - C-Farm Dec 17 v1.xls

G33 F

Inputs and Simulation Control Interface for C-Farm Model

Weatherfile Name and Location: C:\ARK\Research\C-Farm\Pullman.xls Revision: 081111.01

Simulation Start Year: 1963 | Simulation End Year: 1966 | Export to File | Import Settings | Clear All Sheets | Find Weather File | Run C-FARM

Number of Years in the Rotation: 1

	Year 1	Year 2	Year 3	Year 4	Year 5	Year 6	Year 7	Year 8	Year 9	Year 10
Crop 1	Winter Wh.	No Planting								
Crop 2										

Instructions: Crops used **must** be described. Enter "No Planting Event" for years **without** planting events. Perennial Crops will be harvested and killed 3 weeks before next crop planting unless listed earlier with a "Kill Crop" tillage operation.

Describe Up to 5 Crops in Rotation: Winter Wh. | Only describe a crop once

Average Seeding Date	290	Calendar Day (1-365)
Average Date of 50% Flowering	165	Calendar Day (1-365)
Average Maturity Date	200	Calendar Day (1-365)
Maximum Soil Coverage	100	100% = Full Coverage (0-100)
Maximum Rooting Depth	1.99	Lowest depth roots can reach (meters)
Average Expected Yield	3000	kg/ha
Maximum Expected Yield	4000	kg/ha
Minimum Expected Yield	2000	(0-100)
Percent of Moisture in Commercial Yield	12.0	(0-100)
Percent Standing Residue at Harvest	50.0	(0-100)
Percent of Residue Removed	0.0	(0-100)

Harvest Timing as a Percentage of Maturity: (0-100) | Perennials only

Use hourly water infiltration: | Needed for estimation of N2O losses

Daily Crop Outputs: | Last Imported From: | Last Exported To: C:\ARK\Research\C-Farm\EEMAC Copy of C-FARM Nov 11 v1 daily.xls

Daily Soil Outputs: | Last Import Date: Friday, November 28, 2008 | Time: 9:49:29 AM

Daily Weather Outputs: | Last Export Date: Friday, November 28, 2008 | Time: 9:47:58 AM

Annual Soil (Profile) Outputs: | Last Simulation Date: Wednesday, December 17, 2008 | Time: 11:30:19 AM

Season Outputs:

Copyright © 2006-2008. TAMU / WSU. All rights reserved.

Management Control Interface

Sample of Output – Annual summary per soil layer

Soil Carbon Balance							(based on computed yield)				
Year & Layer #	Layer Thickness m	Clay %	Composite Decomposition Factor (unitsless)	Annual Apparent Soil Carbon Decomposition Rate 1/year	Initial Soil Carbon Mg/ha	Apparent Soil Carbon Decomposition Mg C / year	Carbon Input per Layer Mg C / year	Humified Carbon per Layer Mg C / year	Humification Coefficient	Annual Soil Carbon Balance Mg C / ha	
1963	1	0.05	22	0.060	0.003	6.05	0.017	0.000	0.00	-0.017	
1963	2	0.05	22	0.102	0.005	6.05	0.039	0.000	0.00	-0.031	
1963	3	0.13	22	0.199	0.008	15.65	0.118	0.000	0.00	-0.118	
1963	4	0.10	22	0.198	0.004	5.18	0.021	0.000	0.00	-0.021	
1963	5	0.10	22	0.191	0.003	4.02	0.014	0.000	0.00	-0.014	
1963	6	0.10	22	0.183	0.003	2.63	0.007	0.000	0.00	-0.007	
1963	7	0.10	22	0.178	0.002	1.85	0.004	0.000	0.00	-0.004	
1963	8	0.10	22	0.171	0.002	2.37	0.014	0.000	0.00	-0.004	
1963	9	0.50	22	0.163	0.001	3.29	0.004	0.000	0.00	-0.004	
1963	10	0.50	22	0.155	0.001	2.19	0.002	0.000	0.00	-0.002	
1963	11	0.50	22	0.150	0.001	1.43	0.001	0.000	0.00	-0.001	
1963	12	0.50	22	0.147	0.001	1.22	0.001	0.000	0.00	-0.001	
1964	1	0.05	22	0.113	0.005	6.20	0.032	1.855	0.295	0.00	0.265
1964	2	0.05	22	0.133	0.006	6.19	0.038	1.094	0.157	0.00	0.116
1964	3	0.13	22	0.108	0.004	15.90	0.066	1.555	0.226	0.00	0.160
1964	4	0.10	22	0.047	0.001	5.18	0.005	0.052	0.005	0.00	0.000
1964	5	0.10	22	0.046	0.001	4.02	0.003	0.030	0.003	0.00	0.000
1964	6	0.10	22	0.044	0.001	2.63	0.002	0.015	0.001	0.00	0.000
1964	7	0.10	22	0.044	0.001	1.85	0.001	0.010	0.001	0.00	0.000
1964	8	0.10	22	0.041	0.001	2.37	0.013	0.017	0.001	0.00	0.000
1964	9	0.50	22	0.048	0.000	3.29	0.001	0.027	0.003	0.00	0.001
1964	10	0.50	22	0.127	0.001	2.30	0.002	0.100	0.011	0.00	0.009
1964	11	0.50	22	0.122	0.001	1.44	0.001	0.098	0.010	0.00	0.010
1964	12	0.50	22	0.120	0.001	1.22	0.001	0.000	0.000	0.00	-0.001
1965	1	0.05	22	0.085	0.004	6.31	0.005	1.761	0.266	0.00	0.241
1965	2	0.05	22	0.107	0.003	6.39	0.020	0.744	0.105	0.00	0.086
1965	3	0.13	22	0.080	0.002	16.07	0.032	0.567	0.085	0.00	0.052
1965	4	0.10	22	0.048	0.001	5.20	0.005	0.176	0.018	0.00	0.013
1965	5	0.10	22	0.047	0.001	4.02	0.003	0.105	0.011	0.00	0.008
1965	6	0.10	22	0.046	0.001	2.64	0.002	0.067	0.007	0.00	0.005
1965	7	0.10	22	0.046	0.001	1.86	0.001	0.044	0.005	0.00	0.004
1965	8	0.20	22	0.046	0.000	2.28	0.001	0.057	0.006	0.00	0.005
1965	9	0.50	22	0.045	0.000	3.30	0.001	0.069	0.009	0.00	0.000
1965	10	0.50	22	0.062	0.000	2.20	0.001	0.052	0.006	0.00	0.005
1965	11	0.50	22	0.156	0.001	1.45	0.001	0.112	0.013	0.00	0.012
1965	12	0.50	22	0.154	0.001	1.22	0.001	0.000	0.000	0.00	-0.001
1966	1	0.05	22	0.060	0.003	6.39	0.022	1.156	0.178	0.00	0.156
1966	2	0.05	22	0.088	0.005	6.40	0.020	1.340	0.201	0.00	0.170
1966	3	0.13	22	0.030	0.001	16.96	0.019	1.74	0.182	0.00	0.053
1966	4	0.10	22	0.029	0.001	5.21	0.003	0.169	0.018	0.00	0.015
1966	5	0.10	22	0.027	0.000	4.03	0.002	0.096	0.010	0.00	0.008
1966	6	0.10	22	0.024	0.000	2.64	0.001	0.066	0.006	0.00	0.000
1966	7	0.10	22	0.024	0.000	1.85	0.001	0.044	0.005	0.00	0.000
1966	8	0.20	22	0.024	0.000	2.28	0.001	0.057	0.006	0.00	0.005
1966	9	0.50	22	0.023	0.000	3.30	0.001	0.069	0.009	0.00	0.000
1966	10	0.50	22	0.042	0.000	2.20	0.001	0.052	0.006	0.00	0.005
1966	11	0.50	22	0.149	0.001	1.45	0.001	0.112	0.013	0.00	0.012
1966	12	0.50	22	0.147	0.001	1.22	0.001	0.000	0.000	0.00	-0.001
1967	1	0.05	22	0.060	0.003	6.39	0.022	1.156	0.178	0.00	0.156
1967	2	0.05	22	0.088	0.005	6.40	0.020	1.340	0.201	0.00	0.170
1967	3	0.13	22	0.030	0.001	16.96	0.019	1.74	0.182	0.00	0.053
1967	4	0.10	22	0.029	0.001	5.21	0.003	0.169	0.018	0.00	0.015
1967	5	0.10	22	0.027	0.000	4.03	0.002	0.096	0.010	0.00	0.008
1967	6	0.10	22	0.024	0.000	2.64	0.001	0.066	0.006	0.00	0.000
1967	7	0.10	22	0.024	0.000	1.85	0.001	0.044	0.005	0.00	0.000
1967	8	0.20	22	0.024	0.000	2.28	0.001	0.057	0.006	0.00	0.005
1967	9	0.50	22	0.023	0.000	3.30	0.001	0.069	0.009	0.00	0.000
1967	10	0.50	22	0.042	0.001	2.20	0.001	0.052	0.006	0.00	0.005
1967	11	0.50	22	0.149	0.001	1.45	0.001	0.112	0.013	0.00	0.012
1967	12	0.50	22	0.147	0.001	1.22	0.001	0.000	0.000	0.00	-0.001
1968	1	0.05	22	0.060	0.003	6.39	0.022	1.156	0.178	0.00	0.156
1968	2	0.05	22	0.088	0.005	6.40	0.020	1.340	0.201	0.00	0.170
1968	3	0.13	22	0.030	0.001	16.96	0.019	1.74	0.182	0.00	0.053
1968	4	0.10	22	0.029	0.001	5.21	0.003	0.169	0.018	0.00	0.015
1968	5	0.10	22	0.027	0.000	4.03	0.002	0.096	0.010	0.00	0.008
1968	6	0.10	22	0.024	0.000	2.64	0.001	0.066	0.006	0.00	0.000
1968	7	0.10	22	0.024	0.000	1.85	0.001	0.044	0.005	0.00	0.000
1968	8	0.20	22	0.024	0.000	2.28	0.001	0.057	0.006	0.00	0.005
1968	9	0.50	22	0.023	0.000	3.30	0.001	0.069	0.009	0.00	0.000
1968	10	0.50	22	0.042	0.001	2.20	0.001	0.052	0.006	0.00	0.005
1968	11	0.50	22	0.149	0.001	1.45	0.001	0.112	0.013	0.00	0.012
1968	12	0.50	22	0.147	0.001	1.22	0.001	0.000	0.000	0.00	-0.001
1969	1	0.05	22	0.060	0.003	6.39	0.022	1.156	0.178	0.00	0.156
1969	2	0.05	22	0.088	0.005	6.40	0.020	1.340	0.201	0.00	0.170
1969	3	0.13	22	0.030	0.001	16.96	0.019	1.74	0.182	0.00	0.053
1969	4	0.10	22	0.029	0.001	5.21	0.003	0.169	0.018	0.00	0.015
1969	5	0.10	22	0.027	0.000	4.03	0.002	0.096	0.010	0.00	0.008
1969	6	0.10	22	0.024	0.000	2.64	0.001	0.066	0.006	0.00	0.000
1969	7	0.10	22	0.024	0.000	1.85	0.001	0.044	0.005	0.00	0.000
1969	8	0.20	22	0.024	0.000	2.28	0.001	0.057	0.006	0.00	0.005
1969	9	0.50	22	0.023	0.000	3.30	0.001	0.069	0.009	0.00	0.000
1969	10	0.50	22	0.042	0.001	2.20	0.001	0.052	0.006	0.00	0.005
1969	11	0.50	22	0.149	0.001	1.45	0.001	0.112	0.013	0.00	0.012
1969	12	0.50	22	0.147	0.001	1.22	0.001	0.000	0.000	0.00	-0.001
1970	1	0.05	22	0.060	0.003	6.39	0.022	1.156	0.178	0.00	0.156
1970	2	0.05	22	0.088	0.005	6.40	0.020	1.340	0.201	0.00	0.170
1970	3	0.13	22	0.030	0.001	16.96	0.019	1.74	0.182	0.00	0.053
1970	4	0.10	22	0.029	0.001	5.21	0.003	0.169	0.018	0.00	0.015
1970	5	0.10	22	0.027	0.000	4.03	0.002	0.096	0.010	0.00	0.008
1970	6	0.10	22	0.024	0.000	2.64	0.001	0.066	0.006	0.00	0.000
1970	7	0.10	22	0.024	0.000	1.85	0.001	0.044	0.005	0.00	0.000
1970	8	0.20	22	0.024	0.000	2.28	0.001	0.057	0.006	0.00	0.005
1970	9	0.50	22	0.023	0.000	3.30	0.001	0.069	0.009	0.00	0.000
1970	10	0.50	22	0.042	0.001	2.20	0.001	0.052	0.006	0.00	0.005
1970	11	0.50	22	0.149	0.001	1.45	0.001	0.112	0.013	0.00	0.012
1970	12	0.50	22	0.147	0.001	1.22	0.001	0			